Frequency Dependence of ATS-1 Observed Micropulsations on Some Geophysical Parameters

by

Freddie Mason and W. D. Cummings
Physics Department
Grambling College
Grambling, Louisiana 71245

and

P. J. Coleman, Jr.
Department of Planetary and
Space Science
UCLA
Los Angeles, California 90024

This paper is the revised text of SM-50 delivered at the 53rd annual meeting of the American Geophysical Union on April 19, 1972.

May, 1972

Frequency Dependence of ATS-1 Observed Micropulsations on Some Geophysical Parameters

by

Freddie Mason and W. D. Cummings
Physics Department
Grambling College
Grambling, La. 71245

and

P. J. Coleman, Jr.
Dept. of Planetary and Space Science
UCLA
Los Angeles, Calif. 90024

Abstract

Frequency Dependence of ATS-1 Observed Micropulsations on Some Geophysical Parameters. Low frequency oscillations in the earth's magnetic field at the synchronous orbit have been observed with the UCLA magnetometer experiment on board the ATS-1 satellite since Dec. 1966. Some general characteristics of oscillations in the range $2 \times 10^{-3} < f < 20 \times 10^{-3}$ Hz have been previously reported by these authors. A further analysis of oscillations in the above mentioned range for the two year interval Dec. 1966 through Dec. 1968 is reported here. It was found that the frequency of an event increases with the sum of Kp for 24 hours prior to the event midpoint. For events with duration greater than 6 hours, the product moment correlation coefficient for frequency, f, and $\sum$Kp was $R = 0.87$. The best least squares linear fit to the data for these events was $f = (0.32)\sum Kp + 5.1$. For the same events the correlation coefficient for frequency and Dst was $R = -0.83$.

In this paper we will discuss the correlation between the frequency of magnetic oscillations observed at ATS-1 and two indices of geomagnetic disturbance. The indices are $\sum$Kp and $<$Dst$>$. In tabulating $\sum$Kp we summed the Kp index for 24 hours prior to the midpoint of the event as observed at ATS-1. In tabulating $<$Dst$>$ for an event we also computed the average over 24 hours prior to the event midpoint.

We found that the frequency of oscillations is negatively correlated with $\langle Dst \rangle$, i.e., when $\langle Dst \rangle$ is large negative, the event frequency is high. We also found that the frequency of oscillations is well correlated with ΣKp, i.e., when ΣKp is high, the event frequency is also high. When the event frequencies are plotted against ΣKp, it appears that the oscillations can be divided into two groups; those that occur when the plasmapause is beyond the orbit of the satellite, and those that occur when the plasmapause is within the orbit of the satellite.

Slide 1. This slide shows equatorial Dst, taken from the compilation of Sugiura and Poras for the magnetic storm that began on September 20, 1967. On the same plot we have placed the low-frequency oscillations that occurred at ATS-1 during this storm. The occurrence of low-frequency oscillations is marked with a cross on this plot. The horizontal bar shows the duration of the event. The vertical bar is one standard deviation on either side of the average of the measured frequencies for the event. This plot further illustrates the tendency of an event frequency to be high during magnetically disturbed conditions and to decline as the magnetosphere grows progressively quieter during the recovery phase of the storm.

Slide 2. Noting the tendency illustrated in the previous slide, we decided to quantitatively compare the event frequency with $\langle Dst \rangle$ for 24 hours prior to the event midpoint. This slide represents the results of such a comparison. In this plot only those events with duration greater than or equal to 5 hours were selected. The ordinate of the plot gives the event frequency in milli-hertz and the abscissa gives the average value of $\langle Dst \rangle$ for 24 hours prior to a given event. The error bars give one

standard deviation on either side of the average frequency of the event. The solid line is the least squares linear fit to the data. The product moment correlation coefficient for this data was -0.84. We then computed correlation coefficients for frequency and $\langle Dst \rangle$ over 12 hours.

Slide 3. In this plot we have included only those events that had duration greater than or equal to 6 hours. The absolute value of the correlation coefficient was -0.694. As events with shorter duration were included, we found that the correlation coefficient was reduced in absolute value from the value just mentioned.

Realizing that both Dst and Kp decrease together, we decided to quantatively compare the event frequencies with the sum of Kp for 24 hours prior to the event midpoint. The next slide shows the result of one such comparison.

Slide 4. The ordinate of the plot gives the event frequency in milli-hertz and the abscissa is the sum of Kp for 24 hours prior to the event midpoint. In this plot we have included those events greater than or equal to 6 hours. The error bars are of the same nature as those in previous slides. The correlation coefficient was 0.87.

Even though the linear fit to the data had a relatively high correlation coefficient, it appears that the oscillations occurred in two groups; this could be interpretated alternatively. One group occurs when Σ Kp was approximately above 17. The next slide shows our interpretation of this result.

Slide 5. In this plot we have included all events with duration greater than or equal to 5 hours. We calculated linear fits and correlation coeffi-

cients for the two groups of data points. The linear fit for the points with $\Sigma Kp < 17$ had an associated correlation coefficient 0.79. The correlation coefficient for the points with $\Sigma Kp > 17$ was only 0.26, therefore the slope of this line is perhaps not very significant.

We suggest that the dividing line represents the plasmapause; i.e., when $\Sigma Kp < 17$ (or average $Kp < 2$) the plasmapause is beyond $L = 6.5$ on the day side of the magnetosphere, when $\Sigma Kp > 17$ (or average $Kp > 2$) the plasmapause is within $L = 6.5$. We believe this interpretation is consistent with direct satellite measurements of the plasmapause position (particularly the measurements reported by Taylor, Brinton, and Pharo).

In summary: We find that the frequency of magnetic oscillations observed at ATS-1 are well correlated with the two indices of geomagnetic disturbance, $< Dst >$ and ΣKp. Also it appears that the Kp for 24 hours prior to the midpoint ordered the data in such a way that the events occurred in two distinct groups. For $\Sigma Kp <$ about 17 the oscillations occur at about 7 milli-Hz and for $\Sigma Kp >$ about 17 the oscillations occur at about 14 milli-hertz.

We believe that the lower frequency oscillation occur whenever the satellite is within the plasmapause and the higher frequency oscillation occur whenever the satellite is outside the plasmapause.

Acknowledgements

The work done at Grambling College was supported by the National Aeronautics and Space Administration grant NGR-19-011-007. The work done at UCLA was supported by the National Aeronautics and Space Administration grant NGL-05-007-004.


Low-Frequency
Oscillations, ATS-1
and
Equatorial Dst
Dst
20
0
-20
-40
-60
-80
-100
-120
Freq.
(10⁻³ Hz)
16
14
12
10
8
6
4
2
0
19
20
21
22
23
24
25
26
27
28
SEPTEMBER (1967)

Slide 1


Freq.
(10^-3 Hz)
16
14
12
10
8
6
4
2
0
Events with Duration
≥ 5 Hours
-50
-40
-30
-20
-10
0
<Dst>
24 Hours

Slide 2


Freq.
(10^-3 Hz)
16
14
12
10
8
6
4
2
0
ATS - 1
Events with Duration
≥ 6 Hours
-50
-40
-30
-20
-10
0
< Dst >
12 Hours

Slide 3


Freq.
(10⁻³ Hz)
0
2
4
6
8
10
12
14
16
5
10
15
20
25
30
35
40
45
Σ Kp
24 Hours
ATS-1
Events with Duration
≥ 6 Hours

Slide 4


Freq.
(10⁻³ Hz)
0
2
4
6
8
10
12
14
16
0
5
10
15
20
25
30
35
40
45
Σ Kp
24 Hours
ATS-1
Events with Duration
≥ 5 Hours
Plasmapause

Slide 5